


Crop Report

Report name: Loxton Crop report
 Report date: 29/07/2010
 Last climate date available: 28/07/2010
 Client name: MSF
 Paddock name: Loxton Dune
 Report generated by: MSF
 Date sown: 02-Jun
 Crop type: Wheat
 Variety sown: Yitpi
 Sowing density: 120 plants/m²


Weather station used: Loxton
 Rainfall records used: Weather station
 Soil type: Loxton-Low SSC
 Maximum rooting depth: 180 cm
 Roots constrained by EC: yes
 Stubble type: wheat
 Stubble amount: 250 kg/ha
 Start of growing season: 01-Apr
 Initial conditions date: 21-Apr
 Growing season rainfall to date: 97.3 mm
 Date of last rainfall entry: ?
 Expected harvest date: 26-Nov

Grain Yield Outcome




This graph shows the probability of exceeding a range of yield outcomes this season. It takes into account your pre-season soil moisture, the weather conditions so far, soil N and agronomic inputs. The long term record from your nominated weather station is then used to simulate what would have happened from this date on in each of the past 100 years. The yield results are used to produce this graph.

Hay Yield Outcome



This graph shows the probability of exceeding a range of hay yield outcomes this season. It takes into account the same factors as the grain yield graph above. When above ground dry matter is below 2t/ha, hay yield is assumed to be 70% of dry matter, with a moisture content of 13%. When dry matter is between 2 and 12t/ha, hay yield is assumed to be between 70 and 75% of dry matter (sliding scale). When dry matter is above 12t/ha, hay yield is assumed to be between 75 and 80% (sliding scale).


Current dry matter: 68 kg/ha



Predicted									
Earliest	31-Aug	3-Sep	5-Sep	13-Sep	17-Sep	25-Sep	3-Oct	10-Oct	25-Oct
Median	5-Sep	7-Sep	10-Sep	20-Sep	24-Sep	1-Oct	9-Oct	18-Oct	2-Nov
Latest	10-Sep	12-Sep	14-Sep	26-Sep	1-Oct	9-Oct	18-Oct	28-Oct	12-Nov

<i>Percentage of years in which frost occurs</i>		<i>Percentage of years in which heat shock occurs during grain fill (Z70-79)</i>	
Mild		Mild	
Minimum temperature between 2 and 0°C during flowering (Z60-69)	1%	Maximum temperature between 32 and 34°C	68%
Moderate		Moderate	
Minimum temperature between 0 and -2°C during flowering and early grain fill (Z60-75)	0%	Maximum temperature between 34 and 36°C	48%
Severe		Severe	
Minimum temperature less than -2°C during flowering and grain fill (Z60-79)	0%	Maximum temperature above 36°	38%
<i>Incidence of frost for this growing season</i>		<i>Incidence of heat shock for this growing season, during grain fill (Z70-79)</i>	
Mild		Mild	
Minimum temperature between 2 and 0°C during flowering (Z60-69)	0	Maximum temperature between 32 and 34°C	0
Moderate		Moderate	
Minimum temperature between 0 and -2°C during flowering and early grain fill (Z60-75)	0	Maximum temperature between 34 and 36°C	0
Severe		Severe	
Minimum temperature less than -2°C during flowering and grain fill (Z60-79)	0	Maximum temperature above 36°	0

Current distribution of PAW




PAW = Plant Available Water
 CLL = Crop Lower Limit or Wilting Point
 DUL = Drained Upper Limit or Field Capacity
 PAWC = Plant Available Water Capacity
 Current Crop PAW = Soil water currently accessible to the roots down to the current rooting depth
 Soil PAW = Total accessible soil water in the soil profile

Water Budget

Initial PAW status @21-Apr	35 mm
Rainfall since 21-Apr	86 mm
Irigations	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
Evaporation since 21-Apr	53 mm
Transpiration since 21-Apr	1 mm
Deep drainage since 21-Apr	0 mm
Run-off since 21-Apr	0 mm
Current PAW status:	67 mm


Probability of Future Waterlogging Events




Nitrogen Budget

Initial N status @21-Apr	111 kg/ha
Mineralisation since 21-Apr	0 kg/ha
N applications	2-Jun: 4 kg/ha
	: kg/ha
	: kg/ha
	: kg/ha
	: kg/ha
Total N in plant	4 kg/ha
De-nitrification since 21-Apr	0 kg/ha
Leaching	0 kg/ha
Current N status:	113 kg/ha


Current distribution of soil nitrogen (kg/ha)




Availability of Water to Growing Roots




Soil Nitrogen



Water Stress



Nitrogen Stress




Brief periods of mild to moderate stress do not necessarily lead to reduced yield. To see the likely impacts of additional nitrogen fertiliser rates use the Nitrogen and Nitrogen Profit reports.

Mean projected crop performance and requirements for the next 10 days assuming no rain and no added fertiliser.

Date	Growth Stage	Evap (mm)	Daily water use (mm)	Daily N use (kg/ha)	Water available to roots above stress threshold (mm)	Water available to roots above crop lower limit (mm)	N available to roots (kg/ha)
29-Jul	12.8	0.0	0.1	0.5	4.0	7.9	14.4
30-Jul	12.9	0.0	0.1	0.5	4.3	8.2	14.6
31-Jul	13.0	0.0	0.1	0.6	4.4	8.5	14.8
1-Aug	13.1	0.2	0.1	0.6	4.7	9.0	15.0
2-Aug	13.2	0.1	0.1	0.7	5.0	9.5	16.0
3-Aug	13.3	0.2	0.2	0.8	5.3	10.0	17.0
4-Aug	13.4	0.2	0.2	0.8	5.5	10.4	17.9
5-Aug	13.5	0.3	0.2	0.9	5.9	11.0	18.6
6-Aug	13.7	0.1	0.2	1.0	6.4	11.7	19.3
7-Aug	13.8	0.3	0.3	1.1	7.0	12.4	19.7


The water available to roots above the stress threshold is the amount of PAW (mm) above one third of the total water holding capacity of this soil. If the water values are below this stress threshold the water available to roots above the stress threshold will be negative.

The season so far - Growing Season Rainfall Deciles



How much rainfall can I expect?


The Bureau of Meteorology Forecast for the next 3 months



National Seasonal Rainfall Outlook: probabilities July to September 2010

Issued by the bureau of Meteorology 23rd June 2010


How much rainfall can I expect?
 The SOI seasonal forecast for the next 3 months.



The SOI is an index that compares the atmospheric pressure between Tahiti and Darwin. SOI Phases are determined by comparing average monthly SOI values of the past two months. Phases of the SOI have been shown to be related to rainfall variability in a range of locations in Australia and around the world.

	Rainfall
Driest	0 to 48 mm
Drier	48 to 66 mm
Average	66 to 84 mm
Wetter	84 to 116 mm
Wettest	116 to 218 mm

How do seasonal forecasts affect yield expectations?



The 30 day mean SOI for June was 1.31, in May it was 10.48.

Yield outcomes of the current SOI Phase ARE NOT significantly different from yield outcomes of all years. Significance is determined on a 90% probability threshold. (PValue=0.137)

Disclaimer: Yield Prophet information is used entirely at your own risk. You accept all risks and responsibility for losses, damages, costs and other consequences of using Yield Prophet information and reports. To the maximum extent permitted by law, Agricultural Production Systems Research Unit and Birchip Cropping Group excludes all responsibility and liability to any person arising directly or indirectly from using the information generated by Yield Prophet."