


Crop Report

Report name: Natya Crop report
 Report date: 12/08/2010
 Last climate date available: 11/08/2010
 Client name: MSF
 Paddock name: Natya Swale
 Report generated by: MSF
 Date sown: 15-May
 Crop type: Wheat
 Variety sown: Yitpi
 Sowing density: 120 plants/m²


Weather station used: Annuello
 Rainfall records used: Weather station
 Soil type: Sandy Clay Loam (generic)
 Maximum rooting depth: 60 cm
 Roots constrained by EC: no
 Stubble type: wheat
 Stubble amount: 250 kg/ha
 Start of growing season: 01-Apr
 Initial conditions date: 05-May
 Growing season rainfall to date: 135.4 mm
 Date of last rainfall entry: ?
 Expected harvest date: 7-Nov

Grain Yield Outcome



This graph shows the probability of exceeding a range of yield outcomes this season. It takes into account your pre-season soil moisture; the weather conditions so far; soil N and agronomic inputs. The long term record from your nominated weather station is then used to simulate what would have happened from this date on in each of the past 100 years. The yield results are used to produce this graph.


Hay Yield Outcome



This graph show the probability of exceeding a range of hay yield outcomes this season. It takes into account the same factors as the grain yield graph above. When above ground dry matter is below 2t/ha, hay yield is assumed to be 70% of dry matter, with a moisture content of 13%. When dry matter is between 2 and 12t/ha, hay yield is assumed to be between 70 and 75% of dry matter (sliding scale). When dry matter is above 12t/ha, hay yield is assumed to be between 75 and 80% (sliding scale).

Current dry matter: 2638 kg/ha


The season so far - Growing Season Rainfall Deciles





Predicted

Earliest	23-May	6-Jun	17-Jun	24-Jun	6-Jul	17-Jul
Median	23-May	6-Jun	17-Jun	24-Jun	6-Jul	17-Jul
Latest	23-May	6-Jun	17-Jun	24-Jun	6-Jul	17-Jul




Predicted

Earliest	2-Aug	5-Aug	10-Aug	21-Aug	24-Aug	31-Aug	10-Sep	18-Sep	6-Oct
Median	2-Aug	5-Aug	10-Aug	23-Aug	28-Aug	4-Sep	15-Sep	25-Sep	11-Oct
Latest	2-Aug	5-Aug	10-Aug	25-Aug	31-Aug	8-Sep	19-Sep	2-Oct	20-Oct

<i>Percentage of years in which frost occurs</i>	
Mild	
Minimum temperature between 2 and 0°C during flowering (Z60-69)	16%
Moderate	
Minimum temperature between 0 and -2°C during flowering and early grain fill (Z60-75)	1%
Severe	
Minimum temperature less than -2°C during flowering and grain fill (Z60-79)	0%
<i>Incidence of frost for this growing season</i>	
Mild	
Minimum temperature between 2 and 0°C during flowering (Z60-69)	0
Moderate	
Minimum temperature between 0 and -2°C during flowering and early grain fill (Z60-75)	0
Severe	
Minimum temperature less than -2°C during flowering and grain fill (Z60-79)	0

<i>Percentage of years in which heat shock occurs during grain fill (Z70-79)</i>	
Mild	
Maximum temperature between 32 and 34°C	43%
Moderate	
Maximum temperature between 34 and 36°C	16%
Severe	
Maximum temperature above 36°	9%
<i>Incidence of heat shock for this growing season, during grain fill (Z70-79)</i>	
Mild	
Maximum temperature between 32 and 34°C	0
Moderate	
Maximum temperature between 34 and 36°C	0
Severe	
Maximum temperature above 36°	0

Current distribution of PAW




PAW = Plant Available Water
 CLL = Crop Lower Limit or Wilting Point
 DUL = Drained Upper Limit or Field Capacity
 PAWC = Plant Available Water Capacity
 Current Crop PAW = Soil water currently accessible to the roots down to the current rooting depth
 Soil PAW = Total accessible soil water in the soil profile

Current root depth = 594 mm
 Median final root depth = 700 mm
 Current crop PAW = 40 mm
 Soil PAW = 51 mm
 PAWC = 167 mm

Water Budget

Initial PAW status @ 5-May	41 mm
Rainfall since 5-May	101.8 mm
Irrigations	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
Evaporation since 5-May	60 mm
Transpiration since 5-May	29 mm
Deep drainage since 5-May	0 mm
Run-off since 5-May	0 mm
Current PAW status:	51 mm


Probability of Future Waterlogging Events







Nitrogen Budget

Initial N status @ 05-May	188 kg/ha
Mineralisation since 05-May	2 kg/ha
N applications	: kg/ha
	: kg/ha
	: kg/ha
	: kg/ha
	: kg/ha
	: kg/ha
	: kg/ha
Total N in plant	88 kg/ha
De-nitrification since 05-May	0 kg/ha
Leaching	0 kg/ha
Current N status:	103 kg/ha

Current distribution of soil nitrogen (kg/ha)





Brief periods of mild to moderate stress do not necessarily lead to reduced yield. To see the likely impacts of additional nitrogen fertiliser rates use the Nitrogen and Nitrogen Profit reports.

Mean projected crop performance and requirements for the next 10 days assuming no rain and no added fertiliser.

Date	Growth Stage	Evap (mm)	Daily water use (mm)	Daily N use (kg/ha)	Water available to roots above stress threshold (mm)	Water available to roots above crop lower limit (mm)	N available to roots (kg/ha)
12-Aug	33.3	0.5	1.4	1.0	17.8	41.5	28.0
13-Aug	33.7	0.5	1.5	1.0	16.1	40.0	27.3
14-Aug	34.1	0.5	1.5	0.9	14.8	38.6	26.6
15-Aug	34.4	0.5	1.5	0.8	13.6	37.5	26.1
16-Aug	34.8	0.5	1.6	0.7	12.0	35.9	25.5
17-Aug	35.2	0.5	1.6	0.7	10.3	34.2	25.1
18-Aug	35.5	0.6	1.5	0.6	8.8	32.8	24.6
19-Aug	36.0	0.5	1.5	0.6	7.3	31.4	24.2
20-Aug	36.4	0.6	1.4	0.5	6.1	30.3	23.9
21-Aug	36.8	0.6	1.4	0.5	6.3	30.4	23.5

The water available to roots above the stress threshold is the amount of PAW (mm) above one third of the total water holding capacity of this soil. If the water values are below this stress threshold the water available to roots above the stress threshold will be negative.

SOI Phase and analogue years


The SOI is currently in the Rising phase. The 31 day mean SOI for July was 18.74, in June it was 1.31.

The years in history with the same SOI phase:

1898, 1903, 1906, 1912, 1916, 1926, 1928, 1933, 1936, 1939, 1943, 1947, 1948, 1949, 1954, 1960, 1963, 1974, 1979, 1984, 1985, 1988, 1995, 1999, 2003

How much rainfall can I expect?


The SOI seasonal forecast for the next 3 months.



The SOI is an index that compares the atmospheric pressure between Tahiti and Darwin. SOI Phases are determined by comparing average monthly SOI values of the past two months. Phases of the SOI have been shown to be related to rainfall variability in a range of locations in Australia and around the world.

	Rainfall
Driest	0 to 41 mm
Drier	41 to 60 mm
Average	60 to 89 mm
Wetter	89 to 121 mm
Wettest	121 to 235 mm

How do seasonal forecasts affect yield expectations?



The 31 day mean SOI for July was 18.74, in June it was 1.31.

Yield outcomes of the current SOI Phase ARE NOT significantly different from yield outcomes of all years. Significance is determined on a 90% probability threshold. (PValue=0.357)

Disclaimer: Yield Prophet information is used entirely at your own risk. You accept all risks and responsibility for losses, damages, costs and other consequences of using Yield Prophet information and reports. To the maximum extent permitted by law, Agricultural Production Systems Research Unit and Birchip Cropping Group excludes all responsibility and liability to any person arising directly or indirectly from using the information generated by Yield Prophet."