


Crop Report

Report name: Ouyen Crop report
 Report date: 11/11/2010
 Last climate date available: 9/11/2010
 Client name: MSF
 Paddock name: Ouyen Dune
 Report generated by: MSF
 Date sown: 20-May
 Crop type: Wheat
 Variety sown: Yitpi
 Sowing density: 120 plants/m2


Weather station used: Ouyen (Post Office)
 Rainfall records used: Weather station
 Soil type: Dune Loamy Sand (Walpeup)
 Maximum rooting depth: 180 cm
 Roots constrained by EC: yes
 Stubble type: None
 Stubble amount: kg/ha
 Start of growing season: 01-Apr
 Initial conditions date: 12-May
 Growing season rainfall to date: 271.3 mm
 Date of last rainfall entry: ?
 Expected harvest date: 16-Nov

Grain Yield Outcome



This graph shows the probability of exceeding a range of yield outcomes this season. It takes into account your pre-season soil moisture; the weather conditions so far; soil N and agronomic inputs. The long term record from your nominated weather station is then used to simulate what would have happened from this date on in each of the past 100 years. The yield results are used to produce this graph.


Hay Yield Outcome



This graph show the probability of exceeding a range of hay yield outcomes this season. It takes into account the same factors as the grain yield graph above. When above ground dry matter is below 2t/ha, hay yield is assumed to be 70% of dry matter, with a moisture content of 13%. When dry matter is between 2 and 12t/ha, hay yield is assumed to be between 70 and 75% of dry matter (sliding scale). When dry matter is above 12t/ha, hay yield is assumed to be between 75 and 80% (sliding scale).

Current dry matter: 8816 kg/ha

The season so far - Growing Season Rainfall Deciles





Predicted

Earliest	1-Jun	13-Jun	23-Jun	3-Jul	14-Jul	24-Jul
Median	1-Jun	13-Jun	23-Jun	3-Jul	14-Jul	24-Jul
Latest	1-Jun	13-Jun	23-Jun	3-Jul	14-Jul	24-Jul




Predicted

Earliest	12-Aug	15-Aug	18-Aug	1-Sep	5-Sep	13-Sep	23-Sep	4-Oct	20-Oct
Median	12-Aug	15-Aug	19-Aug	1-Sep	5-Sep	13-Sep	24-Sep	4-Oct	20-Oct
Latest	12-Aug	15-Aug	19-Aug	1-Sep	5-Sep	13-Sep	24-Sep	4-Oct	20-Oct

<i>Percentage of years in which frost occurs</i>	
Mild Minimum temperature between 2 and 0°C during flowering (Z60-69)	100%
Moderate Minimum temperature between 0 and -2°C during flowering and early grain fill (Z60-75)	0%
Severe Minimum temperature less than -2°C during flowering and grain fill (Z60-79)	0%
<i>Incidence of frost for this growing season</i>	
Mild Minimum temperature between 2 and 0°C during flowering (Z60-69)	1
Moderate Minimum temperature between 0 and -2°C during flowering and early grain fill (Z60-75)	0
Severe Minimum temperature less than -2°C during flowering and grain fill (Z60-79)	0

<i>Percentage of years in which heat shock occurs during grain fill (Z70-79)</i>	
Mild Maximum temperature between 32 and 34°C	100%
Moderate Maximum temperature between 34 and 36°C	0%
Severe Maximum temperature above 36°	0%
<i>Incidence of heat shock for this growing season, during grain fill (Z70-79)</i>	
Mild Maximum temperature between 32 and 34°C	1
Moderate Maximum temperature between 34 and 36°C	0
Severe Maximum temperature above 36°	0


Current distribution of PAW



Water Budget

Initial PAW status @ 12-May	46 mm
Rainfall since 12-May	239.3 mm
Irrigations	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
Evaporation since 12-May	115 mm
Transpiration since 12-May	133 mm
Deep drainage since 12-May	0 mm
Run-off since 12-May	0 mm
Current PAW status:	39 mm


Probability of Future Waterlogging Events




Nitrogen Budget

Initial N status @ 12-May	83 kg/ha
Mineralisation since 12-May	2 kg/ha
N applications	20-May: 10 kg/ha
	: kg/ha
	: kg/ha
	: kg/ha
	: kg/ha
	: kg/ha
Total N in plant	73 kg/ha
De-nitrification since 12-May	0 kg/ha
Leaching	0 kg/ha
Current N status:	22 kg/ha


Current distribution of soil nitrogen (kg/ha)




Availability of Water to Growing Roots




Soil Nitrogen



Water Stress



Nitrogen Stress



Brief periods of mild to moderate stress do not necessarily lead to reduced yield. To see the likely impacts of additional nitrogen fertiliser rates use the Nitrogen and Nitrogen Profit reports.


Mean projected crop performance and requirements for the next 10 days assuming no rain and no added fertiliser.

Date	Growth Stage	Evap (mm)	Daily water use (mm)	Daily N use (kg/ha)	Water available to roots above stress threshold (mm)	Water available to roots above crop lower limit (mm)	N available to roots (kg/ha)
11-Nov	86.6	0.7	0.5	0.1	-5.9	34.3	21.7
12-Nov	87.0	0.3	0.5	0.1	-6.7	33.5	21.7
13-Nov	89.0	0.2	0.5	0.0	-7.2	33.0	21.7
14-Nov	90.0	0.2	0.3	0.0	-7.6	32.3	21.7
15-Nov	9.0	0.2	0.0	0.0	0.0	0.0	0.0
16-Nov	9.0	0.2	0.0	0.0	0.0	0.0	0.0
17-Nov	9.0	0.2	0.0	0.0	0.0	0.0	0.0
18-Nov	9.0	0.4	0.0	0.0	0.0	0.0	0.0
19-Nov	9.0	0.3	0.0	0.0	0.0	0.0	0.0
20-Nov	9.0	0.7	0.0	0.0	0.0	0.0	0.0

The water available to roots above the stress threshold is the amount of PAW (mm) above one third of the total water holding capacity of this soil. If the water values are below this stress threshold the water available to roots above the stress threshold will be negative.

How much rainfall can I expect?

The Bureau of Meteorology Forecast for the next 3 months




National Seasonal Rainfall Outlook: probabilities October to December 2010

Issued by the bureau of Meteorology 23rd September 2010

Queensland Department of Environment and Resource Management (DERM) 3 month rainfall forecast based on the current phase of the SOI

Probability of exceeding Median Rainfall



SOI Phase and analogue years


The SOI is currently in the Positive phase. The 30 day mean SOI for September was 25.8, in August it was 17.0.

The years in history with the same SOI phase:

1892, 1893, 1894, 1906, 1909, 1910, 1916, 1917, 1920, 1924, 1928, 1938, 1942, 1943, 1945, 1947, 1950, 1954, 1955, 1956, 1960, 1962, 1964, 1967, 1971, 1973, 1974, 1975, 1981, 1988, 1996, 1998, 2000, 2008

How much rainfall can I expect?


The SOI seasonal forecast for the next 3 months.



The SOI is an index that compares the atmospheric pressure between Tahiti and Darwin. SOI Phases are determined by comparing average monthly SOI values of the past two months. Phases of the SOI have been shown to be related to rainfall variability in a range of locations in Australia and around the world.

	Rainfall
Driest	0 to 30 mm
Drier	30 to 48 mm
Average	48 to 71 mm
Wetter	71 to 103 mm
Wettertest	103 to 230 mm

How do seasonal forecasts affect yield expectations?



The 30 day mean SOI for September was 25.8, in August it was 17.0.

Yield outcomes of the current SOI Phase ARE NOT significantly different from yield outcomes of all years. Significance is determined on a 90% probability threshold. (PValue=0.601)

Disclaimer: Yield Prophet information is used entirely at your own risk. You accept all risks and responsibility for losses, damages, costs and other consequences of using Yield Prophet information and reports. To the maximum extent permitted by law, Agricultural Production Systems Research Unit and Birchip Cropping Group excludes all responsibility and liability to any person arising directly or indirectly from using the information generated by Yield Prophet."