


# Crop Report

Report name: Paringa Crop report  
 Report date: 29/07/2010  
 Last climate date available: 28/07/2010  
 Client name: MSF  
 Paddock name: Paringa Dune  
 Report generated by: MSF  
 Date sown: 06-May  
 Crop type: Wheat  
 Variety sown: Gladus  
 Sowing density: 120 plants/m<sup>2</sup>


Weather station used: Meringur (Vic.)  
 Rainfall records used: Weather station  
 Soil type: Carwarp-Moderate SSC  
 Maximum rooting depth: 180 cm  
 Roots constrained by EC: yes  
 Stubble type: wheat  
 Stubble amount: 250 kg/ha  
 Start of growing season: 01-Apr  
 Initial conditions date: 22-Apr  
 Growing season rainfall to date: 69.2 mm  
 Date of last rainfall entry: ?  
 Expected harvest date: 9-Nov

## Grain Yield Outcome



This graph shows the probability of exceeding a range of yield outcomes this season. It takes into account your pre-season soil moisture, the weather conditions so far, soil N and agronomic inputs. The long term record from your nominated weather station is then used to simulate what would have happened from this date on in each of the past 100 years. The yield results are used to produce this graph.

## Hay Yield Outcome




This graph shows the probability of exceeding a range of hay yield outcomes this season. It takes into account the same factors as the grain yield graph above. When above ground dry matter is below 2t/ha, hay yield is assumed to be 70% of dry matter, with a moisture content of 13%. When dry matter is between 2 and 12t/ha, hay yield is assumed to be between 70 and 75% of dry matter (sliding scale). When dry matter is above 12t/ha, hay yield is assumed to be between 75 and 80% (sliding scale).

Current dry matter: 716 kg/ha



**Predicted**

<b>Earliest</b>	1-Jun	13-Jun	22-Jun		1-Jul	14-Jul	23-Jul
<b>Median</b>	1-Jun	13-Jun	22-Jun		1-Jul	14-Jul	23-Jul
<b>Latest</b>	1-Jun	13-Jun	22-Jun		1-Jul	14-Jul	23-Jul




**Predicted**

<b>Earliest</b>	8-Aug	10-Aug	12-Aug	24-Aug	28-Aug	4-Sep	12-Sep	22-Sep	8-Oct
<b>Median</b>	10-Aug	13-Aug	16-Aug	28-Aug	2-Sep	9-Sep	19-Sep	28-Sep	14-Oct
<b>Latest</b>	13-Aug	16-Aug	19-Aug	2-Sep	7-Sep	14-Sep	24-Sep	5-Oct	23-Oct

<p><i>Percentage of years in which frost occurs</i></p> <p><b>Mild</b> Minimum temperature between 2 and 0°C during flowering (Z60-69) 5%</p> <p><b>Moderate</b> Minimum temperature between 0 and -2°C during flowering and early grain fill (Z60-75) 0%</p> <p><b>Severe</b> Minimum temperature less than -2°C during flowering and grain fill (Z60-79) 0%</p>		<p><i>Percentage of years in which heat shock occurs during grain fill (Z70-79)</i></p> <p><b>Mild</b> Maximum temperature between 32 and 34°C 43%</p> <p><b>Moderate</b> Maximum temperature between 34 and 36°C 19%</p> <p><b>Severe</b> Maximum temperature above 36° 13%</p>	
<p><i>Incidence of frost for this growing season</i></p> <p><b>Mild</b> Minimum temperature between 2 and 0°C during flowering (Z60-69) 0</p> <p><b>Moderate</b> Minimum temperature between 0 and -2°C during flowering and early grain fill (Z60-75) 0</p> <p><b>Severe</b> Minimum temperature less than -2°C during flowering and grain fill (Z60-79) 0</p>		<p><i>Incidence of heat shock for this growing season, during grain fill (Z70-79)</i></p> <p><b>Mild</b> Maximum temperature between 32 and 34°C 0</p> <p><b>Moderate</b> Maximum temperature between 34 and 36°C 0</p> <p><b>Severe</b> Maximum temperature above 36° 0</p>	

**Current distribution of PAW**



**PAW** = Plant Available Water  
**CLL** = Crop Lower Limit or Wilting Point  
**DUL** = Drained Upper Limit or Field Capacity  
**PAWC** = Plant Available Water Capacity  
**Current Crop PAW** = Soil water currently accessible to the roots down to the current rooting depth  
**Soil PAW** = Total accessible soil water in the soil profile

**Water Budget**

Initial PAW status @ 22-Apr	22 mm
Rainfall since 22-Apr	76.2 mm
Irigations	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
Evaporation since 22-Apr	57 mm
Transpiration since 22-Apr	8 mm
Deep drainage since 22-Apr	0 mm
Run-off since 22-Apr	0 mm


**Current PAW status: 34 mm**

**Nitrogen Budget**


Initial N status @ 22-Apr	41 kg/ha
Mineralisation since 22-Apr	0 kg/ha
N applications	8-May: 9 kg/ha
	: kg/ha
	: kg/ha
	: kg/ha
	: kg/ha
	: kg/ha
Total N in plant	22 kg/ha
De-nitrification since 22-Apr	0 kg/ha
Leaching	0 kg/ha

**Current N status: 31 kg/ha**


**Probability of Future Waterlogging Events**




**Current distribution of soil nitrogen (kg/ha)**




Availability of Water to Growing Roots




Soil Nitrogen



Water Stress



Nitrogen Stress




Brief periods of mild to moderate stress do not necessarily lead to reduced yield. To see the likely impacts of additional nitrogen fertiliser rates use the Nitrogen and Nitrogen Profit reports.

**Mean projected crop performance and requirements for the next 10 days assuming no rain and no added fertiliser.**

Date	Growth Stage	Evap (mm)	Daily water use (mm)	Daily N use (kg/ha)	Water available to roots above stress threshold (mm)	Water available to roots above crop lower limit (mm)	N available to roots (kg/ha)
29-Jul	16.0	0.2	0.2	0.2	6.1	29.4	18.1
30-Jul	16.0	0.2	0.2	0.2	5.6	29.1	18.0
31-Jul	16.0	0.3	0.2	0.2	5.2	28.9	17.9
1-Aug	16.0	0.3	0.2	0.2	4.8	28.7	17.9
2-Aug	16.0	0.4	0.2	0.2	4.5	28.5	17.8
3-Aug	16.0	0.5	0.2	0.2	4.2	28.3	17.7
4-Aug	16.0	0.5	0.2	0.2	4.2	28.7	17.6
5-Aug	16.0	0.4	0.2	0.2	4.9	29.5	17.5
6-Aug	16.0	0.5	0.2	0.2	4.9	29.5	17.4
7-Aug	16.0	0.6	0.2	0.2	5.5	30.3	17.3


The water available to roots above the stress threshold is the amount of PAW (mm) above one third of the total water holding capacity of this soil. If the water values are below this stress threshold the water available to roots above the stress threshold will be negative.

The season so far - Growing Season Rainfall Deciles



How much rainfall can I expect?


The Bureau of Meteorology Forecast for the next 3 months



National Seasonal Rainfall Outlook: probabilities July to September 2010

Issued by the bureau of Meteorology 23rd June 2010


**How much rainfall can I expect?**  
 The SOI seasonal forecast for the next 3 months.



The SOI is an index that compares the atmospheric pressure between Tahiti and Darwin. SOI Phases are determined by comparing average monthly SOI values of the past two months. Phases of the SOI have been shown to be related to rainfall variability in a range of locations in Australia and around the world.

	Rainfall
Driest	0 to 46 mm
Drier	46 to 68 mm
Average	68 to 87 mm
Wetter	87 to 118 mm
Wettest	118 to 212 mm

**How do seasonal forecasts affect yield expectations?**



The 30 day mean SOI for June was 1.31, in May it was 10.48.

Yield outcomes of the current SOI Phase ARE NOT significantly different from yield outcomes of all years. Significance is determined on a 90% probability threshold. (PValue=0.156)

Disclaimer: Yield Prophet information is used entirely at your own risk. You accept all risks and responsibility for losses, damages, costs and other consequences of using Yield Prophet information and reports. To the maximum extent permitted by law, Agricultural Production Systems Research Unit and Birchip Cropping Group excludes all responsibility and liability to any person arising directly or indirectly from using the information generated by Yield Prophet."